



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 9/16, 47/18		A2	(11) International Publication Number: WO 98/40069
			(43) International Publication Date: 17 September 1998 (17.09.98)
(21) International Application Number: PCT/EP98/01478 (22) International Filing Date: 13 March 1998 (13.03.98) (30) Priority Data: 97104200.7 13 March 1997 (13.03.97) EP (34) Countries for which the regional or international application was filed: DE et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(71) Applicant (for all designated States except US): HEXAL AG [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): KLOKKERS, Karin [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). KUTSCHERA, Marion [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). FISCHER, Wilfried [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). (74) Agents: BOETERS, Hans, D. et al.; Boeters & Bauer, Bereiteranger 15, D-81541 München (DE).		Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: STABILIZATION OF ACID SENSITIVE BENZIMIDAZOLS WITH AMINO/CYCLODEXTRIN COMBINATIONS			
(57) Abstract <p>A pharmaceutical formulation comprising or consisting of: a benzimidazole derivative as active ingredient, and as excipients, at least one cyclodextrin and at least one amino acid.</p>			

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

STABILIZATION OF ACID SENSITIVE BENZIMIDAZOLS WITH AMINO/CYCLODEXTRIN COMBINATIONS

The present invention relates to stable pharmaceutical formulations, containing moisture and acid sensitive benzimidazole derivatives (e. g. omeprazole) as pharmaceutically active ingredient combined with amino acids and cyclodextrins as excipients, and to a method for preparation such pharmaceutical formulations.

Omeprazole (5-methoxy-2-(4-methoxy-3,5-dimethyl-2-pyridinylmethyl-sulfinyl)-1H-benzimidazol) is an effective inhibitor of gastric acid secretion and has a strong antiulcer activity. It is known, that omeprazole rapidly decomposes at acidic and neutral pH. Furthermore moisture, organic solvents and UV-irradiation accelerate the degradation of omeprazole too, causing discoloration of the substance in solution, as well as in solid

form. For example, omeprazole has half-life time of 10 minutes in an aqueous solution of below a pH-value of 4, but 18 hours at pH 6,8 and about 300 days at pH 11 (M. Mathew and co-workers, Drug. Dev. Ind. Pharm., 21,965,1995). The drug has been reported to be stable under alkaline conditions [Pilbrant A° and Cederberg C. Scand. J. Gastroenterology, Suppl. 108, 113-120(1985)]. According to A. Brändström and co-workers (Acta Chem. Scand. 43,536,1989) the acid-catalyzed degradation kinetics of omeprazole is very complicated, the primary degradation is followed by rather complex secondary reactions.

Several methods for stabilizing the acid-unstable compound, in particular omeprazole have been described.

Some patent applications (USP-5232706, EPA-0567201 A2, EPA-0519144 A1, EPA-0496437 A2, USP-5385739, DEA-1204363 and EPA-0247983 B1) claim a common method to overcome this stability problem by applying an inert protective layer between the core and the enteric coating layer. The core contains the pharmaceutical active substance (omeprazole) or its salts, alkaline or acid neutralizing additives, alkaline salts or a combination thereof.

The resorption of omeprazole occurs in the upper duodenum. Therefore, a quick and complete release of the active ingredient after passage of the pylorus must be ensured in order to guarantee a sufficiently high bioavailability. For this, omeprazole is provided with a coating of enteric, i.e. gastric juice-resistant material, which is insoluble in the acid environment of the stomach (ca. pH 1 to 3) on the one hand, but dissolves in the weakly acidic to weakly alkaline region of the duodenum (pH > 5,5). Ordinary enteric coatings, however, are made of acidic compounds. If the core containing omeprazole will be covered with a conventional enteric coating without an subcoating, omeprazole rapidly decomposes by direct or indirect

contact with the coating, with the result that the preparation become discolored.

Although the sensitivity of omeprazole against organic solvent is known, acetone and methylene-chloride (EPA-0496437 A2, EPA-0567201 A2) or acetone and ethanol (USP-5385739, EPA-0519144 A1) are used for the enteric coating of the tablets. This treatment can damage the active ingredient during the enteric coating process or during the long-term storage.

All known procedures consist of complicated multi-step operations and result in expensive final products, which must be stored under specific conditions in moisture proof packages.

DE-427785 A1, DE-3427786 A1, DE-3427787 A1 intended to solve the stability problems of omeprazole by a different method.

Omeprazole and β -cyclodextrin (CD) or derivatives of β -CD (hydroxypropylcyclodextrin) were reacted in 96% ethanol for 15 hours at elevated temperature. Upon cooling a white crystalline substance was isolated, which was believed to be an omeprazole/ β -CD inclusion complex. However the elevated temperature through 15 hours in the presence of 96 % ethanol results in extensive degradation of omeprazole thus there is hardly active ingredient remained in the isolated product. It is generally known, that ethanol is a competing cyclodextrin-complex forming agent. From a 96 % ethanolic system only the crystalline ethanol/ β -CD complex can be isolated by using the mentioned method (Otagiri, M. et al: Acta Pharm. Suetica 21, 357 (1984), Pitha, J. and Hoshino, T.: Int. J. Pharm 80, 234 (1992)).

The WO 93/13138 discloses a method for stabilization of acid-sensitive benzimidazoles, more specifically for the stabilization of omeprazole in drug formulations, which comprise a cyclodextrin-complex of omeprazole, a protective inert layer and an enteric coating. The omeprazole is reacted in presence of alkaline hydroxides, alkaline salts, amines or buffers with

cyclodextrin and derivatives for 1 to 30 minutes at 30 to 70 °C in a homogeneous solution system. After cooling to room temperature the reacted solution is allowed to stand at 4 °C for 3 to 15 hours to form the omeprazole/cyclodextrin-complex. The isolated inclusion-complex is washed with some cooled water several times to completely remove the remaining alkaline component on the inclusion complex. Alternatively, from the reacted solution the water might be removed by spray drying, freeze drying or vacuum evaporating for isolation of the inclusion complex powder as stable compound.

In the state of the art a core made of omeprazole and an alkaline substance as well as a inclusion complex from omeprazole and cyclodextrin without an amino acid is not stable enough. A inert protective layer is necessary to guarantee the stability of omeprazole and specific moisture-proof packages were needed for storing the final product.

Main object of the invention is to guarantee a stabilization of benzimidazoles such as omeprazole as active ingredient by forming a benzimidazole/cyclodextrin inclusion complex.

It has now been found, that benzimidazoles such as omeprazole can be stabilized by complexation with a cyclodextrin such as β -cyclodextrin in the presence of an amino acid. It has further been found that in this case surprisingly no additional inert or enteric layer is needed to protect particles or a core containing the benzimidazole/cyclodextrin complex and an amino acid. Merely optionally the core may be coated directly with an enteric coating layer.

Thus, the problem underlying the invention is solved by a pharmaceutical formulation comprising or consisting of

- a benzimidazole derivative as active ingredient, and as excipients
- at least one cyclodextrin and
- at least one amino acid.

The present invention does provide a new pharmaceutical benzimidazole formulation with improved stability features and simplified preparation process.

The benzimidazole derivative can be a compound which is decomposed in the presence of humidity and especially at a pH ≤ 11 , especially ≤ 7 . Examples for these benzimidazole derivatives are omeprazole, lansoprazole, leminoprazole, rabeprazole, and pantoprazole. Omeprazole is preferred.

Further, a specific embodiment of the invention concerns a pharmaceutical formulation, wherein the inclusion complex forming agent is β -cyclodextrin, mono- or polyalkylated β -cyclodextrin, mono- or polyhydroxyalkylated β -cyclodextrin or γ -cyclodextrin, preferably β -cyclodextrin.

The amino acid useful for the pharmaceutical formulation according to the invention can be an alkaline amino acid, preferably arginine, lysine or hydroxy lysine and especially L-arginine, L-lysine or L-hydroxy lysine; an alkaline dipeptide or a pharmaceutically acceptable alkaline amino acid derivate.

Further, a specific embodiment of the invention concerns a pharmaceutical formulation, wherein the molar ratio of omeprazole to cyclodextrin is 1 to 10 and preferably 1 to 2.

Further, a specific embodiment of the invention concerns a pharmaceutical formulation, wherein the molar ratio of the amino

acid (preferably L-arginine) to omeprazole is 0.5 to 10 and preferably 1 to 1.

Further, a specific embodiment of the invention concerns a pharmaceutical formulation, wherein the formulation is a powdered, pelletized or granulated form, optionally processed to tablets.

The pharmaceutical formulation according to the invention can be characterized in that the particles of the powder, of the granulate or of the pelletized formulation are not coated with an enteric coating. The powder, the granulate or the pelletized formulation can be, however, contained in capsules which can optionally be provided with an enteric coating.

Further, the particles of the powder, of the granulate or of the pelletized formulation can be provided with an enteric coating and optionally be contained in capsules which are not provided with an enteric coating.

As examples for enteric coating materials polymeres such as cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, copolymerized methacrylic acid/methacrylic acid methyl esters or water-based polymer dispersions, for instance, compounds known under the trade name Eudragit® L (Röhm Pharma), or similar compounds can be used. The enteric coating layer can optionally contain a pharmaceutically acceptable plasticizer such as dibutylphthalate, diethylsebacat or tri-ethylcitrat. Dispersants such as talc, colorants and pigments may also be included to the enteric coating layer.

The problem underlying the invention is, in addition, solved by a process for the production of a pharmaceutical composition according to the invention, wherein

- (i) a benzimidazole derivative, at least one cyclodextrin, and at least one amino acid are wetted with water and mixed;
- (ii) the resulting mixture is dried.

Further, the problem underlying the invention is solved by a process for the production of a pharmaceutical composition according to the invention, wherein

- (i) a benzimidazole derivative, at least one cyclodextrin, and at least one amino acid are wetted with water and mixed;
- (ii) the resulting mixture is dried; and
- (iii) discoloration of the composition is examined and if a discolored product is obtained, said discolored product is discarded, another amino acid is selected and steps (i) to (iii) are repeated until an uncolored product is obtained.

The mixing in step (i) of the process according to the invention can be carried out by wet-kneading.

The water to be used in step (i) of the process according to the invention can be ammoniacal water or can be free of any ammonia.

The drying in step (ii) of the process according to the invention can be carried out by freeze-drying, spray-drying or vacuum-drying.

For the production of pellets the pharmaceutical formulation according to the invention can be mixed with a binding agent such as microcrystalline cellulose and an excipient such as hydroxypropylcellulose and moistened, for example with isopropanol, and then formulated into pellets by conventional pharmaceutical procedures. The pellets can be used as cores for

further processing. The pellets may be filled directly into capsules which are optionally coated with an enteric coating. Further, the pellets themselves may be coated with an enteric coating and optionally filled into uncoated capsules.

The enteric coating layer can be applied onto the pellets by conventional coating techniques such as, for instance, pan coating, fluidized bed coating, fluidized bed bottom sprayed coating or a Turbo Jet-Technology for the production of large amounts using dispersions of polymers in water and/or suitable organic solvents or by using latex suspensions of said polymers. Examples for enteric coating polymers have already been mentioned.

An application of the pharmaceutical formulation according to the invention results in pharmaceutically effective plasma levels and offers a sufficiently high bioavailability. This could not be expected in view of the fact that the active ingredient is used in combination with a complexing agent.

In order to describe the invention more specifically but without intending to limit the scope of the invention in any way the following examples are presented:

Example 1 and comparative examples

Compositions containing omeprazole, β -cyclodextrin and an amino acid at a molar ratio of 1:2:1 were prepared by kneading in presence of water and powdered after drying. Cellulose acetate phthalate, an acidic reacting excipient, was mixed to the composition in an amount of 5 w/w %, calculated to the total weight of the sample. The optical density of the composition powders after storing for 7 days at 60 °C in presence of 96% R.H. is illustrated in Table I.

Table I.: Composition and discoloration (optical density measured at 346 nm)of omeprazole + β -cyclodextrin mixtures, in presence of amino acids and cellulose acetate phthalate after 7 days at 60 °C and 96% relative humidity

Sample	omeprazole	β -CD	amino acid	cellulose acetate phthalate	O.D. after dissolving the powders
A	+	+			1.0
B	+	+		+	2.4
C	+	+	arginine		0.4
D	+	+	arginine	+	0.8
E	+	+	lysine		0.6
F	+	+	lysine	+	0.7

The presence of an amino acid enhances the stability of the inclusion complex of omeprazole and β -cyclodextrin as illustrated in Table I. There is no rapid decomposition of omeprazole by direct contact with cellulose acetate phthalate under stressed conditions.

Comparative Example

Inclusion complexes of omeprazole and β -cyclodextrin were prepared by the same method as described before but without using an amino acid.

As reference omeprazole and lactose mixtures were prepared, with similar weight-ratios. The molar ratio of omeprazole to β -cyclodextrin and to lactose was 1:2. The result is illustrated in Table II.

Table II: Composition and discoloration of powder mixtures stored at 40 °C at 76% R.H. for 20 days

samples	omeprazole	β -CD	lactose	cellulose acetate phthalate	stored in closed container	stored in open container	O.D. after dissolving the powders
G	+		+		+		0,2
H	+		+	+	+		0,4
I	+	+			+		0,2
J	+	+		+	+		0,4
K	+		+			+	0,8
L	+		+	+		+	2,4
M	+	+				+	0,7
N	+	+		+		+	2,0

Stability of the inclusion complex in absence of an amino acid seems to be acceptable only by storage in closed containers even in the absence of cellulose acetate phthalate. The pre-sence of cellulose acetate phthalate in all cases enhances the degradation of omeprazole. Comparing the samples stored in closed and in open containers the role of the humidity is quite obvious: the discoloration of omeprazole in open containers is much higher in all cases than in the closed containers. The degradation is significantly accelerated by humidity (samples stored in open containers) and by the presence of cellulose acetate phthalate (acidic additive), the β -cyclodextrin itself is not a significantly better stabilizer than the lactose.

Example 2

In further experiments the β -cyclodextrin has been suspended in diluted aqueous ammonium hydroxide solution, before omeprazole and arginine has been added. The samples were prepared as described before and stored at 50 °C and 76% R.H. for 7 days. Cellulose acetate phthalate (CAP) (5%w/w) was mixed to all samples after the β -cyclodextrin/omeprazole/arginine amino acid suspensions were dried and powdered. The composition of the samples as well as their discoloration are shown in Table III.

Table III.: Excipients added to omeprazole and cellulose acetate phthalate, method of preparation and the discoloration of the samples after storing for 7 days at 50°C and 76% R.H. in open containers

Sample	Components			Methods of preparation	O.D. of the solution
O	lactose		CAP	powder	2.4
P	β -CD		CAP	mixture	1.6
				powder	
				mixture	
Q	β -CD+NH ₃	arginine	CAP	powder	0.6
R	β -CD		CAP	mixture	0.8
				powder	
				mixture	
S	β -CD	arginine	CAP	wet	0.3
T	β -CD+NH ₃	arginine	CAP	kneading	0.1
				wet	
				kneading	
U	lactose	arginine	CAP	wet	1.2
V	β -CD+NH ₃		CAP	kneading	0.9
				wet	
				kneading	

These data compared to the data of Table I. and II. clearly demonstrate, that while β -cyclodextrin - when used in wet kneading or in solution - alone is more effective than lactose in protecting omeprazole against discoloration particularly when it is reacted with omeprazole in ammonia-alkaline solution, its protecting effect is significantly potentiated by the presence of arginine or lysine.

The lactose/arginine combination (U) or the β -cyclodextrin+NH₃ without arginine (V) did not result in satisfactory stabilizing effect. The required omeprazole protecting effect (against acid and water provoked decomposition) could be attained by the ternary combination of omeprazole/ β -CD/arginine (S-T), prepared

by wet kneading in water, wherein the water can be ammonia-alkaline water or water free of ammonia.

During the drying process the ammonia has been completely removed, in the end-product ammonia can not be detected.

Particularly important is that this combination is not sensitive to 76% R.H., at elevated temperature.

Example 3

208 g L-arginine are dissolved in 2L distilled water, and 400 g omeprazole are suspended in this solution (Suspension I).

3 kg β -cyclodextrin (water content 11.95%) are suspended with 3.2 L distilled water by Ultra-Turrax for 5 minutes (Suspension II.).

Suspension I is poured into Suspension II. under vigorous stirring by Ultra-Turrax for 15 minutes at 8000 rpm.

For isolation of the solid product the suspension is frozen and the water content is removed by freeze-drying.

Yield: 3242 g (97.3%)

omeprazole content: 12.3%

Water content: 2.5%

Determination of omeprazole content of samples

As it is shown in Table IV., the samples showed a good storage stability. The decrease of the omeprazole content in the samples - stored under stressed conditions - does not exceed an absolute value of 0.5%, at samples - stored at ambient tem-

perature - practically no change in active ingredient content was observed.

Visual observation of the samples showed no color change, except of the sample stored in open container at daylight (see Table IV.). The moisture absorption of the samples - stored at 76% RH. - was remarkable, without significant discoloration (Table V.)

Table IV.: Omeprazole content of the samples after two weeks storage under stressed conditions and 6 months storage at ambient temperature

storage conditions	storage period	omeprazole content (% ^a ±SD)		Appearance
		"a"	"b"	
-	-	12.3±0.0 8	12.0±0.10	off white powder
40°C, 76% RH	2 weeks	12.0±0.0 6	11.7±0.05	not changed
ambient temperature - closed container - open container	6 months 6 months	12.3±0.2 3	12.1±0.23 11.3±0.5	not changed very light yellowish color

* related to the dry substance

Table V.: Moisture absorption of the samples

storage conditions	storage period	loss on drying (%)	
		"a"	"b"
-	-	2.79	2.16
40°C, 76% R.H.	2 weeks	9.17	8.23
ambient temperature - closed container - open container	6 months 6 months	2.65 -	2.37 4.35

Example 4

0.64 g omeprazole and 5.08 g β -cyclodextrin (water content: 12%) are homogenized in a mortar, then a solution of 0.33 g lysine in 1.5 ml of 2.5% NH_3 is added and homogenization is continued. Finally the obtained suspension is granulated through a laboratory sieve with 0.4 mm and dried at 45°C for 24 hours. 5.5 g of granules is obtained.
omeprazole content: 10.9%

Example 5

Sample a:

1.32 g omeprazole, 0.68 g L-arginine, and 10.56 g β -cyclodextrin (water content: 11.9%) were powdered by co-grinding in a ball-mill, then kneaded with 3 ml of water for a few minutes. The resulting paste was dried over P_2O_5 at room temperature in a vacuum exsiccator overnight, ground roughly to granule-size particles.

To characterize the stability of omeprazole in this formulation also the following samples were prepared with and without amino acid and/or β -cyclodextrin (water content: 11.9%):

Sample b (without β -cyclodextrin):	1.32 g omeprazole
	0.68 g L-arginine
	9.3 g lactose

Sample c (without arginine):	1.32 g omeprazole
	10.56 g β -cyclodextrin

Example 7

First three mixtures were prepared:

- 1) 4.1 g omeprazole and 6 g β -cyclodextrin (water content: 11,9 %)
- 2) 25 g β -cyclodextrin and 55 g water
- 3) 21 g water and 2,1 g L-arginine

Then the three mixtures were mixed together and the resulting suspension was spray-dried under the following conditions:

inlet temperature:	120 - 125°C
outlet temperature:	75 - 80°C
air pressure:	2.5 kg/cm ²
feeding speed:	4 ml/min

37.5 g off-white powder is obtained.

omeprazole content: 12.6%

L-arginine content: 6.22%

water content (KFT): 5.40%

Example 8

509 g pharmaceutical formulation (omeprazole: β -cyclodextrin: arginine) (1:2:1), 163 g microcrystalline cellulose and 55 g hydroxypropylcellulose are mixed for 5 minutes. Then 270 g isopropanol are given to the mixture and mixed for 10 minutes on high level. After that the mixture is extruded and instantly worked up to pellets. The pellets are dried for about 16 - 18 hours at 40 °C.

The pellets can be filled into hard gelantine capsules optionally enteric coated. Or the pellets are enteric coated with Eudragit L, for example L 100-55, L100 or L 30D according to standard methods.

Claims

1. A pharmaceutical formulation comprising or consisting of
 - a benzimidazole derivative as active ingredient, and as excipients
 - at least one cyclodextrin and
 - at least one amino acid.
2. Pharmaceutical formulation according to claim 1, in which the benzimidazole derivative is selected from benzimidazole derivatives which are decomposed in the presence of humidity and especially at a pH ≤ 11 , especially ≤ 7 .
3. Pharmaceutical formulation according to any of the preceding claims, in which the benzimidazole derivative is omeprazole.
4. Pharmaceutical formulation according to any of the preceding claims, in which the cyclodextrin is β -cyclodextrin, mono- or polyalkylated β -cyclodextrin, mono- or polyhydroxyalkylated β -cyclodextrin or γ -cyclodextrin.
5. Pharmaceutical formulation according to any of the preceding claims, in which the cyclodextrin is β -cyclodextrin.
6. Pharmaceutical formulation according to any of the preceding claims, in which the amino acid is a basic amino acid, preferably arginine, lysine or hydroxy lysine, especially L-arginine, L-lysine or L-hydroxy lysine; an alkaline dipeptide or a pharmaceutically acceptable alkaline amino acid derivative.

7. Pharmaceutical formulation according to any of the preceding claims, characterized by a molar ratio of omeprazole to cyclodextrin of from 1 to 10 and preferably 1 to 2.
8. Pharmaceutical formulation according to any of the preceding claims, characterized by a molar ratio of amino acid (preferably L-arginine) to omeprazole of from 0,5 to 10 and preferably 1 to 1.
9. Pharmaceutical formulation according to any of the preceding claims in powdered, granulated or pelletized form, each optionally processed to tablets.
10. Pharmaceutical formulation according to claim 9, characterized in that the particles of the powder, of the granulate or of the pelletized formulation are not coated with an enteric coating.
11. Pharmaceutical formulation according to claim 10, characterized in that the powder, the granulate or the pelletized formulation is contained in a capsule which is optionally provided with an enteric coating.
12. Pharmaceutical formulation according to claim 9, characterized in that the particles of the powder, of the granulate or of the pelletized formulation are provided with an enteric coating and optionally contained in a capsule which is not provided with an enteric coating.
13. Process for the production of a pharmaceutic composition according to any of the preceding claims, characterized in that
 - (i) a benzimidazole derivative, at least one cyclodextrin and at least one amino acid are

wetted with water and mixed;

(ii) the resulting mixture is dried.

14. Process for the production of a pharmaceutical composition according to any of the preceding claims, characterized in that

(i) a benzimidazole derivative, at least one cyclodextrin and at least one amino acid are wetted with water and mixed;

(ii) the resulting mixture is dried; and

(iii) discoloration of the composition is examined and if a discolored product is obtained, said discolored product is discarded, another amino acid is selected and steps (i) to (iii) are repeated until an uncolored product is obtained.

15. Process according to claim 13 or 14, characterized in that in step (i) of claim 13 the mixing is carried out by wet-kneading.

16. Process according to any of claims 13 to 15, characterized in that in step (i) of claim 13 the water is ammoniacal water or is free of ammonia.

17. Process according to any of claims 13 to 16, characterized in that in step (ii) of claim 13 the drying is carried out by freeze-drying, spray-drying or vacuum-drying.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/44, 9/16, 47/18, 47/48	A3	(11) International Publication Number: WO 98/40069 (43) International Publication Date: 17 September 1998 (17.09.98)
(21) International Application Number: PCT/EP98/01478 (22) International Filing Date: 13 March 1998 (13.03.98) (30) Priority Data: 97104200.7 13 March 1997 (13.03.97) EP (34) Countries for which the regional or international application was filed: DE et al. (71) Applicant (for all designated States except US): HEXAL AG [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): KLOKKERS, Karin [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). KUTSCHERA, Marion [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). FISCHER, Wilfried [DE/DE]; Industriestrasse 25, D-83607 Holzkirchen (DE). (74) Agents: BOETERS, Hans, D. et al.; Boeters & Bauer, Bereiteranger 15, D-81541 München (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 17 December 1998 (17.12.98)
(54) Title: STABILIZATION OF ACID SENSITIVE BENZIMIDAZOLES WITH AMINO ACID/CYCLODEXTRIN COMBINATIONS (57) Abstract A pharmaceutical formulation comprising or consisting of: a benzimidazole derivative as active ingredient, and as excipients, at least one cyclodextrin and at least one amino acid.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/01478

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/44 A61K9/16 A61K47/18 A61K47/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	CHEMICAL ABSTRACTS, vol. 124, no. 16, 15 April 1996 Columbus, Ohio, US; abstract no. 211738, XP002079802 see abstract & S.J. HWANG ET AL.: "A COMPARATIVE STUDY ON THE PHARMACEUTICAL PROPERTIES OF RECTAL SUPPOSITORY CONTAINING OMEPRAZOLE COMPLEXES" YAKCHE HAKHOECHI, vol. 25, no. 3, 1995, pages 227-237, ---	1-17
Y	EP 0 444 625 A (HANMI PHARM. IND. CO. LTD., KR) 4 September 1991 see the whole document ---	1-17
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

7 October 1998

Date of mailing of the international search report

16/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Scarponi, U

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/01478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DE 34 27 787 A (BYK GULDEN LOMBERG) 30 January 1986 cited in the application see the whole document ---	1-17
Y	DE 34 27 786 A (BYK GULDEN LOMBERG) 30 January 1986 cited in the application see the whole document ---	1-17
Y	WO 96 24338 A (ASTRA) 15 August 1996 see the whole document ---	1-17
Y	US 5 232 706 A (A. PALOMO COLL) 3 August 1993 see the whole document ---	1-17
Y	WO 96 38175 A (TAKEDA) 5 December 1996 see the whole document ---	1-17
Y	WO 93 13138 A (SUNKYONG IND. CO. LTD.,KR) 8 July 1993 cited in the application see the whole document ---	1-17
Y	CHEMICAL ABSTRACTS, vol. 119, no. 22, 29 November 1993 Columbus, Ohio, US; abstract no. 234039, XP002079803 see abstract & JP 05 194225 A (YOSHITOMI) 3 August 1993 ---	1-17
Y	DATABASE WPI Week 9332 Derwent Publications Ltd., London, GB; AN 93-255964 '32! XP002079804 see abstract & KR 9 208 161 B (HANMI PHARM. IND. CO.,KR) 24 September 1992 ---	1-17
A	WO 94 02140 A (YOSHITOMI) 3 February 1994 see the whole document -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/01478

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 444625	A	04-09-1991	CA 2037101 A,C DE 69102307 D DE 69102307 T ES 2057628 T JP 2038521 C JP 4234817 A JP 7051503 B US 5219870 A	28-08-1991 14-07-1994 09-02-1995 16-10-1994 28-03-1996 24-08-1992 05-06-1995 15-06-1993
DE 3427787	A	30-01-1986	AU 4636885 A WO 8600913 A EP 0190239 A	25-02-1986 13-02-1986 13-08-1986
DE 3427786	A	30-01-1986	NONE	
WO 9624338	A	15-08-1996	AU 695774 B AU 4683996 A BR 9605121 A CA 2186037 A CN 1146720 A CZ 9602918 A EP 0752851 A FI 964028 A HU 9603112 A JP 9511768 T NO 964271 A PL 316685 A SK 126196 A	20-08-1998 27-08-1996 14-10-1997 15-08-1996 02-04-1997 16-07-1997 15-01-1997 08-10-1996 28-08-1997 25-11-1997 08-10-1996 03-02-1997 08-10-1997
US 5232706	A	03-08-1993	NONE	
WO 9638175	A	05-12-1996	AU 5780696 A JP 9048730 A	18-12-1996 18-02-1997
WO 9313138	A	08-07-1993	BR 9207000 A CA 2127111 A,C CN 1076124 A,B DE 69222950 D DE 69222950 T EG 20115 A	28-11-1995 08-07-1993 15-09-1993 04-12-1997 28-05-1998 31-07-1997

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/01478

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9313138 A		EP 0619825 A	19-10-1994
		ES 2111148 T	01-03-1998
		HU 70494 A	30-10-1995
		JP 2662518 B	15-10-1997
		JP 7506088 T	06-07-1995
		KR 9608231 B	21-06-1996
		MX 9207676 A	01-06-1993
		US 5399700 A	21-03-1995
WO 9402140 A	03-02-1994	AU 4513393 A	14-02-1994
		JP 7033659 A	03-02-1995

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.